

Factors affecting the fetal hemoglobin levels in patients with sickle cell anemia and thalassemia

Citation for published version (APA):

Dimovski, A. J. (1993). *Factors affecting the fetal hemoglobin levels in patients with sickle cell anemia and thalassemia*. [Doctoral Thesis, Maastricht University]. Rijksuniversiteit Limburg.
<https://doi.org/10.26481/dis.19931125ad>

Document status and date:

Published: 01/01/1993

DOI:

[10.26481/dis.19931125ad](https://doi.org/10.26481/dis.19931125ad)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 05 May. 2023

Stellingen

behorende bij het proefschrift

Factors affecting the fetal hemoglobin levels
in patients with
sickle cell anemia and thalassemia

van

Aleksandar Jovo Dimovski

Maastricht
25 november 1993

1. The polymorphic patterns of the $(AT)_x(T)_y$ repeats at -530 5' to the β -globin gene do not correlate with the level of Hb F in patients with β -thalassemia major.
2. The determination of specific sequence variations in the 5' flanking region of the G_γ - and A_γ -globin genes is a reliable, time- and cost-effective procedure for the determination of the major β^S haplotypes.
3. The C \rightarrow T substitution at position -158 5' to the G_γ gene is the most consistent sequence variation associated with the β^S and β -thal chromosomes which are capable of conferring high Hb F expression under the hematopoietic stress of these diseases.
4. The possible influence of the sequence variations in the 5' flanking region of the G_γ - and A_γ -globin genes and in the LCR HS-2 on the overall expression of the γ -globin genes should be tested in an experimental system where the conditions of hematopoietic stress could be reproduced, i.e. transgenic mice.
5. The silencing effect of the 3' HS-1 sequences should be further evaluated through studies of patients with additional deletions or mutations within this sequence.
6. The improvement in DNA methodology, particularly the PCR-related procedures, makes the determination of various Hb variants by this technique a method of choice when compared to the classical protein chemistry analysis.

7. Gene therapy in hemoglobinopathies is hampered mainly by the low efficiency of injection, the inability to obtain sufficiently enriched stem cell population, and the risks of insertional mutagenesis.
8. The improvement in early survival of patients with acute myocardial infarction treated with streptokinase has still not been proven to be associated with a long-term survival benefit.
9. The determination of the frequencies of particular DNA polymorphisms should be established in each population prior to their implementation in the forensic medical practice.
10. Administration of iron derivatives in addition to erythropoietin facilitates the correction of anemia in patients with normal and even increased iron stores.
11. Mild mutations of the CFTR gene can be the cause of infertility in otherwise normal male individuals.
12. Molecular medicine should be included as a separate discipline in the curriculum of the Medical Faculties.
13. The European Economic Community should recognize The Republic of Macedonia under its genuine name.